

# **Genetic diversity in *Neospora caninum***

**By**

**SARWAT EKRAM AL-QASSAB**

**A thesis submitted in fulfillment of the requirements for the  
degree of Doctor of Philosophy**



**Department of Medical and Molecular Biosciences**

**Faculty of Science**

**University of Technology, Sydney**

**Australia**

**2009**

## **Certificate**

This study was conducted in the Department of Medical and Molecular Biosciences, Faculty of Science, University of Technology Sydney under the supervision of Professor John Ellis and Adjunct Professor Michael Reichel.

I certify that the work in this thesis has not previously been submitted for a degree nor has it been submitted as part of requirement for a degree except as fully acknowledged within the text.

I also certify that the thesis has written by me and help that I received in my research work and the preparation of the thesis itself has been acknowledged. In addition, I also certify that the thesis has been written by me with additional help from Prof. John Ellis and Prof. Michael Reichel as acknowledged in individual chapters. Furthermore I certify that all information sources and literatures used are indicated in the thesis.

**SARWAT EKRAM AL-QASSAB**

## Acknowledgment

My deepest gratitude goes to my parents. Their unconditional support and blessings made it possible for me to carry on my studies. My greatest debt of gratitude and appreciation is to my wife Shahnaz, whose help, courage and support are behind my achievement. I appreciate her forbearance and patience during the course of my research. I feel obliged to mention my children Israa, Eman and Mustafa who have shown a great sense of responsibilities, understanding and courage.

I would like to express my respect and sincere gratitude to my supervisor Professor John Ellis, for his expert guidance, constructive criticism and continuous support and encouragement throughout my study. It has been a privilege to work with him. His assistance will always be remembered.

I would like to thank my co-supervisor Professor Michael Reichel for his valuable suggestion and comments throughout the period of my candidature.

Special thanks to Dr Najah Nassif for her technical guidance. Her advices are deeply appreciated.

I would like to thank this University for giving me the opportunity to conduct my study and I also appreciated their financial support.

I would like to thank all researchers mentioned through this thesis for the donation of parasite strains and DNA for conduct of this study.

Many thank to Peter Williams (Kemps Creek farm) for the donation of dog puppies for isolation of *Toxoplasma gondii*.

Special thanks to Dr Debra Birch, for her valuable assistance on electron microscopy of *T. gondii*.

I am also grateful to Dr Mary Davey, Professor Peter Windsor, Dr Michael Johnson, Dr Craig Hall, and my fellow colleagues from the lab, Amanda Hudson, Catherine James, Irene Sotirchos, Michael Lees, Robert Walker and Matthew Padula.

## **Refereed publications arising from this thesis**

Al-Qassab, S., Reichel, P. M., Ivens, A., Ellis, J. T. (2009) Genetic diversity detected amongst isolates of *Neospora caninum* and development of a multiplex PCR for typing strains. *Molecular and Cellular Probes*, **23**: 132-9.

Al-Qassab, S., Reichel, M. P., Hall, G., Johnson, M., Windsor, P. A., Jenkins, D., Su, C., Dubey, J. P., Ellis, J. T. (2009) Isolation of *Toxoplasma gondii* from the brain of a dog in Australia and its biological and molecular characterisations. *Veterinary Parasitology*, **Available online**.

Al-Qassab, S., Reichel, M. P., Ellis, J. T. (2009) A second generation Multiplex PCR for typing strains of *Neospora caninum* using six DNA targets. *Molecular and Cellular Probes*, **Available online**.

## **Papers presented at scientific conferences**

Al-Qassab, S., Reichel, M. P., Hall, G., Johnson, M., Windsor, P. A., Jenkins, D., Su, C., Dubey, J. P., Ellis, J. T. (2009) Isolation of *Toxoplasma gondii* from the brain of a dog in Australia and its biological and molecular characterisations. Australian Society of Parasitology Meeting, Adelaide, Australia.

Al-Qassab, S., Reichel, M. P., Ivens, A., Ellis, J. T. (2009) Genetic diversity detected amongst isolates of *Neospora caninum* and development of a multiplex PCR for typing strains. British Society of Parasitology, Edinburgh, Scotland.

## Abbreviations

ITS1	internal transcribed spacer sequences 1
nPCR	nested PCR
PBS	phosphate-buffered saline
PBST	phosphate-buffered saline Tween-20
PI	post inoculation
RT	room temperature
ss-rDNA	small subunit ribosomal DNA

## Abstract

The apicomplexan parasite *Neospora caninum* has been identified as an important cause of disease in the dairy industry because it causes abortion and neonatal mortality in livestock such as cattle, sheep and goats and also causing neuromuscular disorders and death in dogs.

Biological diversity within the species *N. caninum* isolates has not been studied in detail. Limited differences were reported using several genetic approaches such as sequencing of ribosomal DNA. Mini- and microsatellite DNA markers have recently been reported to represent a valuable tool to study genetic diversity within this parasite. This is the topic of this thesis.

In the present study, mini- and microsatellite DNA markers were used successfully to genotype different strains of *N. caninum* which were isolated from bovine and canine hosts and from different geographical regions of the world. Several mini- and microsatellites were identified that detected differences among different isolates of this parasite. A prototype multiplex PCR assay was developed for strain typing of *N. caninum* based on the length polymorphisms detected amongst three different repetitive markers two- minisatellites and one microsatellite. This method is simple, rapid, highly informative and sensitive for the study of *N. caninum* diversity.

Discrimination amongst isolates can be improved by the incorporation of more loci into a multiplex PCR and so additional DNA markers were sought. The study of genomic DNA resulted in the identification of two additional minisatellites and one microsatellite. A second generation multiplex PCR assay was developed for multilocus strain typing of *N. caninum* based on length polymorphisms of the six satellites. This multiplex included three microsatellites and three minisatellites. This assay provides a higher degree of resolution and provides a valuable tool to study genetic variation among different isolates of this parasite.

Various uses of multiplex PCR were investigated in the study of *N. caninum*. Isolation of *N. caninum* from a litter of pups was attempted; the bitch was previously identified as seropositive to *N. caninum*. Multiplex PCR of DNA from the serum of the bitch showed a genotype for *N. caninum* not previously reported in Australia. *Toxoplasma gondii* was isolated from the brain of a dog for the first time in Australia. The identity of the parasite was confirmed by PCR, Western blotting, electron microscopy and cat bioassay. Genotyping of the isolate (TgDgAu1) was determined by

10 PCR-RFLP markers and revealed it to be a Type II strain. Western blotting demonstrated the presence of IgM antibodies to *T. gondii* suggesting the bitch was probably infected during pregnancy and the *T. gondii* was transmitted to the pups congenitally therefore, this represents the first description of a natural case of congenitally transmission of *T. gondii* in the dog. Unfortunately *N. caninum* was not isolated during these studies.

## Table of contents

<b>Certificate</b>	.....	<b>ii</b>
<b>Acknowledgment</b>	.....	<b>iii</b>
<b>Refereed publications arising from this thesis.....</b>		<b>iv</b>
<b>Abriviation</b>	.....	<b>v</b>
<b>Abstract</b>	.....	<b>vi</b>
<b>Table of contents</b>	.....	<b>viii</b>
<b>List of Tables</b>	.....	<b>xiii</b>
<b>List of Figures</b>	.....	<b>xiv</b>
<b>Chapter 1: General Introduction.....</b>		<b>1</b>
1.1. Introduction.....		2
1.2. Host and host range.....		2
1.3. Economic impact.....		3
1.4. Biology of <i>Neospora caninum</i> .....		4
1.4.1. Life cycle.....		4
1.5. Mode of transmission.....		8
1.6. Epidemiology.....		9
1.7. Control Strategies.....		10
1.7.1. Test'n'Cull.....		10
1.7.2. Therapeutic treatment.....		11
1.7.3. Vaccination.....		11
1.7.4. Postnatal risk/exposure.....		11
1.7.5. Embryo transfer.....		11
1.7.6. Do nothing.....		12
1.8. Pathogenesis of infection / abortion.....		12
1.8.1. In cattle.....		12
1.8.2. In dog.....		14
1.8.3. In goat.....		14
1.8.4. In rodent.....		15



1.9.	Differentiation of <i>Neospora caninum</i> from other closely related coccidia.....	15
1.9.1.	Clinical signs of disease.....	15
1.9.2.	Morphology.....	15
1.9.2.A.	Tissue cysts.....	15
1.9.2.B.	Rhoptries.....	16
1.9.2.C.	Micronemes.....	16
1.9.2.D.	Micropores.....	16
1.9.3.	Protein content.....	16
1.9.4.	DNA characteristics.....	16
1.9.5.	Growth rate.....	17
1.9.6.	Biochemistry.....	18
1.10.	Comparison of biological characteristics between <i>Neospora caninum</i> isolates.....	18
1.10.1.	Ultrastructure.....	18
1.10.1.A.	Tachyzoite size.....	18
1.10.1.B.	Rhoptries.....	18
1.10.1.C.	Micronemes.....	19
1.10.1.D.	Tissue cysts.....	19
1.10.1.E.	Oocysts.....	24
1.10.2.	Pathogenicity.....	30
1.10.3.	Antigens.....	31
1.10.4.	DNA analyses.....	32
1.10.4.A.	18S-like rDNA (ss-rDNA).....	32
1.10.4.B.	Internal transcribed spacer sequences (ITS1).....	33
1.10.4.C.	RAPD-PCR.....	33
1.10.4.D.	Microsatellite.....	34
1.10.4.E.	Nc5 DNA region.....	34
1.10.4.F.	$\alpha$ - and $\beta$ -tubulin genes.....	34
1.10.4.G.	Heat shock protein 70 gene.....	34
1.10.4.H.	Surface antigen gene.....	34
1.10.4.I.	P37, IDA8 and Histone 2A genes.....	35
1.10.5.	Growth rate.....	35

1.10.6. Biochemistry.....	36
1.11. Application of polymerase chain reaction in <i>Neospora caninum</i> .....	36
1.11.1. Repetitive sequences.....	36
1.11.2. Genotyping of parasites.....	38
1.11.3. Mini- and microsatellite markers applied to detect genetic diversity within <i>N. caninum</i> .....	39
1.12. Aims.....	39
 <b>Chapter 2: Genetic diversity detected amongst isolates of <i>Neospora caninum</i> and development of a multiplex PCR for typing strains.....</b>	
2.1. Introduction.....	42
2.2. Material and methods.....	43
2.2.1. Isolates of <i>N. caninum</i> .....	43
2.2.2. Genomic DNA preparation.....	46
2.2.3. Identification of mini- and microsatellite sequences in <i>N. caninum</i> DNA.....	46
2.2.4. PCR of repetitive DNA and DNA sequencing.....	46
2.2.5. Multiplex PCR typing of <i>N. caninum</i> isolates.....	52
2.3. Results.....	52
2.3.1. Characterization of repetitive sequences.....	52
2.3.2. Use of multiplex PCR for the characterization of <i>N. caninum</i> isolates.....	54
2.4. Discussion.....	68
 <b>Chapter 3: A second generation multiplex PCR for typing strains of <i>Neospora caninum</i> using six DNA targets.....</b>	
3.1. Introduction.....	72
3.2. Materials and Methods.....	73
3.2.1. Parasites used in this study.....	73

3.2.2. DNA extraction.....	73
3.2.3. Detection of mini- and microsatellite repetitive sequences in <i>N. caninum</i> DNA.....	73
3.2.4. PCR of repetitive DNA and DNA sequencing.....	74
3.2.5. Diversity among <i>N. caninum</i> isolates through distance analyses..	77
3.3. Results.....	77
3.3.1. Identification of repetitive sequences.....	77
3.3.2. Comparison of isolates by PCR and multiplex PCR.....	77
3.3.3. <i>Neospora hughesi</i> and other related coccidian species.....	81
3.4. Discussion.....	95

#### **Chapter 4: Isolation of *Toxoplasma gondii* from the brain of a dog in Australia and its biological and molecular**

<b>characterisation.....</b>	<b>97</b>
4.1. Introduction.....	98
4.2. Material and methods.....	99
4.2.1. Host history.....	99
4.2.2. Preparation of brain and cultivation of the parasite.....	99
4.2.3. Bioassay in mice.....	99
4.2.4. Bioassay in cats.....	100
4.2.5. DNA extraction.....	101
4.2.6. PCR amplification and sequence analysis.....	101
4.2.7. PCR-RFLP genotyping.....	102
4.2.8. Immunoblot analysis.....	102
4.2.9. Electron microscopy.....	104
4.2.10. Characterisation of <i>N. caninum</i> DNA from sera from infected dog.....	104
4.3 Results.....	104
4.3.1. Genotyping infected dog sera with multiplex PCR.....	104
4.3.2. <i>In vitro</i> isolation of the parasite .....	105
4.3.3. Mouse infection .....	105
4.3.4. PCR analysis .....	105

4.3.5. Cat infection .....	106
4.3.6. PCR-RFLP genotyping.....	106
4.3.7. Western blotting .....	106
4.3.8. Electron microscopy .....	106
4.4. Discussion.....	111
 <b>Chapter 5: General Discussion.....</b>	 114
<b>Chapter 6: References.....</b>	<b>121</b>

## List of Table

<b>Table 1.1.</b> Summary of reported <i>Neospora caninum</i> isolates from different hosts.....	<b>20</b>
<b>Table 1.2A.</b> Size of tachyzoites in different <i>Neospora caninum</i> isolates.....	<b>25</b>
<b>Table 1.2B.</b> Number of rhoptries present in different <i>N. caninum</i> tachyzoites.....	<b>25</b>
<b>Table 1.3.</b> Size of tissue cysts from different infected hosts.....	<b>26</b>
<b>Table 1.4.</b> Sizes and numbers of oocysts shed by dogs.....	<b>28</b>
<b>Table 2.1.</b> Isolates of <i>Neospora caninum</i> used in this study.....	<b>44</b>
<b>Table 2.2.</b> A summary of the mini- and microsatellite loci studied and their PCR amplification conditions.....	<b>47</b>
<b>Table 2.3.</b> Repetitive sequences found in different <i>N. caninum</i> isolates following sequencing of PCR products.....	<b>55</b>
<b>Table 3.1.</b> Isolates of <i>Neospora caninum</i> and <i>N. hughesi</i> used in this study with the number of repetitive sequences found in five loci of different isolates.....	<b>75</b>
<b>Table 3.2.</b> Description of mini- and microsatellite loci and amplification conditions.....	<b>79</b>
<b>Table 4.1.</b> Summary of genotyping data for the TgDgAu1 isolate of <i>T. gondii</i>	<b>107</b>

## List of Figures

<b>Fig. 1.1.</b> Shows the Life cycle of <i>Neospora caninum</i> .....	<b>6</b>
<b>Fig. 2.1.</b> PCR amplification of (A) (Tand-4) and (B) (Tand-15).....	<b>57</b>
<b>Fig. 2.2.</b> PCR amplification of repetitive sequences from different <i>N. caninum</i> isolates. (A) Tand-12; (B) Tand-13; (C) Tand-3.....	<b>58</b>
<b>Fig. 2.3.</b> Multiplex PCR was performed for the three markers: Tand-3 Tand-12 and Tand-13.....	<b>58</b>
<b>Fig. 2.4.</b> DNA sequence alignments of repetitive sequences from various <i>N. caninum</i> isolates.....	<b>60</b>
<b>Fig. 2.5.</b> Sensitivity of multiplex PCR analysis of NC-Nowra using different DNA concentrations.....	<b>66</b>
<b>Fig. 2.6.</b> Stability of multiplex PCR analysis using NC-Nowra cultured for various time periods.....	<b>66</b>
<b>Fig. 2.7.</b> Multiplex PCR of DNA extracted from pup brains showing NC-Liverpool isolate is present in the brains.....	<b>66</b>
 <b>Fig.3.1.</b> PCR amplification of repetitive sequences from different <i>N. caninum</i> isolates. (A) Cont-6, (B) Cont-16 and Cont-14.....	 <b>82</b>
<b>Fig. 3.2.</b> Multiplex PCR was performed for the six markers Tand-3, Tand-12, Tand-13, Cont-6, Cont-14 and Cont-16.....	<b>82</b>
<b>Fig. 3.3.</b> DNA sequence alignments of repetitive sequences from various <i>N. caninum</i> isolates.....	<b>84</b>
<b>Fig. 3.4.</b> Multiplex PCR was used for six markers using DNA from related coccidia.....	<b>92</b>
<b>Fig. 3.5.</b> Stability of multiplex PCR analysis using NC-Nowra cultured for various time periods.....	<b>92</b>
<b>Fig. 3.6.</b> Genetic distances amongst <i>N. caninum</i> isolates based on the sequence data of mini- and microsatellites using GenAlEx6 program.....	<b>94</b>
<b>Fig. 4.1.</b> Western blotting using rat anti- <i>T. gondii</i> serum (A) or the female dog serum (B, C) and tachyzoite extracts. Arrows pointing to differences between the two isolates (Dog3 is TgDgAu1). B and C. were developed with anti-IgG (B) or anti-IgM (C).....	<b>108</b>

<b>Fig 4.2.</b> EM images showing rhoptry (R) and nucleus (N). (A) ME49;	
(B) TgDgAu1.....	<b>108</b>
<b>Fig. 4.3.</b> Multiplex PCR of DNA extracted from dog serum that was	
antibody positive to <i>N. caninum</i> .....	<b>110</b>